

Sufficient number of integrals for the p^{th} order Lyness equation

Dinh T. Tran, Peter H van der Kamp and G. R.W Quispel

Department of Mathematics and Statistics, La Trobe University, Victoria 3086,
Australia

E-mail: dinhtran82@yahoo.com

Abstract. We present a sufficient number of first integrals for the Lyness equation of arbitrary order. We first use the staircase method [16] to construct integrals of a derivative equation of the Lyness equation. Closed-form expressions for the integrals are given based on a non-commutative Vieta expansion. The integrals of the Lyness equation then follow directly from these integrals. Previously found integrals for the Lyness equation arise as special cases of our new set of integrals.

PACS numbers: 02.30.Ik

Submitted to: *J. Phys. A: Math. Gen.*

1. Introduction

We consider the p^{th} order Lyness equation or the p -dimensional Lyness mapping

$$u_n u_{n+p} = a + u_{n+1} + u_{n+2} + \dots + u_{n+p-1}. \quad (1)$$

This equation is a generalization of the equation which was first written by Lyness in 1942 [14] where $p = 2$ and $a = 1$. Recently, equation (1) and its particular cases have been studied by several authors [2, 3, 5, 6, 7, 8, 12]. One important aspect in the study of the Lyness equation (1), or difference equations in general, is finding its integrals (invariants). Up to now, only three integrals of the p^{th} order Lyness equation have been found. The first invariant was given in [7, 10, 11]. The second and the third invariants were discovered in 2004 by Gao et al with the help of computer algebra [8], as well as fourth invariants for the 7th and 8th order Lyness equations. The authors conjectured that the p^{th} order Lyness equation has up to $\lfloor (p+1)/2 \rfloor$ integrals, which is a sufficient number of integrals for complete integrability in the sense of Liouville-Arnold (a $2N$ -dimensional symplectic map has N functionally independent integrals which are in involution with respect to the symplectic structure [4, 23]).

In a recent paper [9], Grammaticos et al have pointed out that the Lyness equation (1) satisfies two criteria of integrability namely having singularity confinement and polynomial growth. This means that the Lyness equation is a good candidate for integrability in the sense of Liouville-Arnold. The authors of [9] also bilinearized the following consequence of equation (1),

$$u_{n+p}(1 + u_n) = u_{n+1}(1 + u_{n+p+1}), \quad (2)$$

which we will call the derivative Lyness equation. In [9], it was shown that the associated bilinear equation is a reduction of the integrable Hirota-Miwa equation. However, this does not allow us to calculate first integrals of the derivative equation.

There exists a method, called the staircase method, to construct integrals for a certain class of ordinary difference equations. This method was introduced in [15, 16] and generalised in [21, 22, 18]. Integrals were obtained for ordinary difference equations (ODEs), derived as travelling wave reductions of integrable partial difference equations (PDEs), i.e. PDEs which exhibit a Lax pair. A so called monodromy matrix is constructed by taking a product of Lax matrices (L, M) along a one period segment of the periodic staircase. By expanding the trace of the monodromy matrix in powers of the spectral parameter, one will obtain integrals for the corresponding ODE. To give closed-form expressions for integrals, in [20] we used this method in conjunction with a non-commutative Vieta expansion and a certain way of splitting Lax matrices. In that paper integrals for reductions of ABS equations [1], were expressed in terms of multi-sums of products, Ψ .

In this paper, we observe that the derivative Lyness equation (2) is a travelling wave reduction of a certain two-dimensional integrable PDE. It turns out that therefore one can use the staircase method to obtain integrals for the derivative Lyness equation. Moreover, we will show that the Lax matrix L for equation (2) fits in the framework

of [20], i.e. we obtain the integrals of the derivative Lyness equation in closed-form. Integrals of the p^{th} order Lyness equation (1) are then obtained directly from those of the derivative Lyness equation. This settles the conjecture given in [8] in the affirmative. Furthermore, this result indicates that the Lyness equation is a good candidate for completely integrability in Liouville-Arnold sense.

The paper is organized as follows. In section 2, we will recall the staircase method for $(p, -1)$ -travelling wave reductions. In section 3, we will give a Lax pair for the partial difference equation associated with equation (2). This Lax pair is derived from a Lax pair for equation (26) in [13] and a gauge transformation. Then, we will give closed-form expressions for integrals of the derivative Lyness equation (2). Once again these integrals can be expressed in terms of multi-sums of products, Ψ , introduced in [20]. We will derive integrals of the original equation (1) from the integrals of the derivative Lyness equation in section 5. One will see that all three integrals given in [8] can be derived as special cases from our closed-form expressions. Finally, we give some relations between the integrals and we discuss functional independence of the integrals.

2. Staircase method for $(p, -1)$ -travelling wave reductions

In this section, we recall the staircase method which was introduced in [15, 16]. We consider a 2-dimensional P Δ E with field variable u ,

$$f(u_{l,m}, u_{l+1,m}, u_{l,m+1}, u_{l+1,m+1}; \alpha) = 0, \quad (3)$$

and parameters $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_k)$. A (s_1, s_2) -reduction can be performed to find solutions that satisfy the periodicity condition $u_{l,m} = u_{l+s_1, m+s_2}$ [21, 22, 18]. For our purpose here, we take $s_1 = p$ and $s_2 = -1$. We introduce a similarity variable $n = l+mp$. Then, $u_{l,m}$ satisfies the periodicity $u_{l,m} = u_{l+p, m-1}$, and the P Δ E reduces to the O Δ E

$$f_n = f(u_n, u_{n+1}, u_{n+p}, u_{n+p+1}; \alpha) = 0. \quad (4)$$

We suppose that equation (3) arises as the compatibility condition of two linear equations, that is, it has a Lax pair. A Lax pair $L_{l,m}, M_{l,m}$ for a P Δ E (3) is a pair of matrices that satisfies, cf [17],

$$L_{l,m} M_{l,m}^{-1} - M_{l+1,m}^{-1} L_{l,m+1} = 0, \quad (5)$$

for solutions of the equation. Similarly, an O Δ E has a Lax pair if there are non-singular matrices $\mathcal{L}_n, \mathcal{M}_n$ that satisfy

$$\mathcal{M}_n \mathcal{L}_n - \mathcal{L}_{n+1} \mathcal{M}_n = 0. \quad (6)$$

The monodromy matrix \mathcal{L}_n for the $(p, -1)$ -reduction is given by, cf [16],

$$\mathcal{L}_n = M_n^{-1} \prod_{i=0}^{\widehat{p-1}} L_{i+n}, \quad (7)$$

where the inversely ordered product is

$$\prod_{i=a}^{\widehat{b}} L_i := L_b L_{b-1} \dots L_{a+1} L_a. \quad (8)$$

Taking $\mathcal{M}_n = L_n$, we obtain a Lax pair $\mathcal{L}_n, \mathcal{M}_n$ for the reduced O Δ E (4) from the Lax pair of the corresponding P Δ E (3). This was first observed in [17]. Lax pairs for O Δ Es obtained from general (s_1, s_2) -reductions have been studied in [18].

It follows from equation (6) that the trace of \mathcal{L}_n is invariant under the map obtained from an O Δ E. Since the Lax matrices generally depend on a spectral parameter, integrals for the O Δ E (4) are obtained by expanding the trace of the monodromy matrix (or powers thereof, or its determinant) in powers of the spectral parameter.

3. Lax pair for the derivative Lyness equation

The derivative Lyness equation (2) is an $(p, -1)$ -traveling reduction of the following P Δ E,

$$u_{l,m+1}(1 + u_{l,m}) = u_{l+1,m}(1 + u_{l+1,m+1}). \quad (9)$$

It is easy to see that this equation is equivalent to the equation

$$(u_{l,m} - 1/2)(u_{l+1,m} + 1/2) = (u_{l+1,m+1} - 1/2)(u_{l,m+1} + 1/2), \quad (10)$$

if we use a transformation $u_{l,m} \mapsto -u_{m,l} - 1/2$ and interchange l and m . It is important to note that equation (10) is a special case of equation (31) in [13], which has a Lax pair. Therefore, a Lax pair for the P Δ E associated with the derivative Lyness equation has the following form,

$$L_{l,m} = \begin{pmatrix} \lambda & -u_{l+1,m} \\ 1 & \frac{\lambda u_{l+1,m}}{u_{l+1,m} + 1} \end{pmatrix}, \quad M_{l,m} = \begin{pmatrix} \lambda & -(u_{l,m} + 1)u_{l,m+1} \\ 1 & 0 \end{pmatrix}. \quad (11)$$

Now we will use a gauge transformation to obtain a more suitable form of a Lax pair for this equation. Recall that if a P Δ E has a Lax pair $(L_{l,m}, M_{l,m})$, then a gauge matrix $G_{l,m}$, which can depend on dependent/independent variables and on the spectral parameter, will give us a new Lax pair $(\tilde{L}_{l,m}, \tilde{M}_{l,m})$ where

$$\tilde{L}_{l,m} = G_{l+1,m} L_{l,m} G_{l,m}^{-1}, \quad \tilde{M}_{l,m} = G_{l,m+1} M_{l,m} G_{l,m}^{-1}. \quad (12)$$

Applying the gauge

$$G_{l,m} = \begin{pmatrix} 1/\lambda & 0 \\ 0 & 1 \end{pmatrix}, \quad (13)$$

to the Lax pair $(L_{l,m}/\lambda, M_{l,m}/\lambda)$, we obtain a new Lax pair for equation (9)

$$\tilde{L}_{l,m} = \begin{pmatrix} 1 & \frac{-u_{l+1,m}}{\lambda^2} \\ 1 & \frac{u_{l+1,m}}{u_{l+1,m} + 1} \end{pmatrix}, \quad \tilde{M}_{l,m}^{-1} = \begin{pmatrix} 0 & 1 \\ \frac{-\lambda^2}{u_{l,m+1}(u_{l,m} + 1)} & \frac{\lambda^2}{u_{l,m+1}(u_{l,m} + 1)} \end{pmatrix}. \quad (14)$$

We will show in the next section that this new Lax pair does fit in the framework of [20].

4. Closed-form expressions for integrals of the derivative Lyness equation

In this section, we will give closed-form expressions for integrals of the derivative Lyness equation (2). These integrals again can be expressed in terms of multi-sums of products, Ψ [20]. We also calculate the integrating factors of these integrals. Recall that $\Lambda_n(u_n, u_{n+1}, \dots, u_{n+p+1})$ is an integrating factor corresponding to an integral $I_n = I(u_n, u_{n+1}, \dots, u_{n+p})$ of the equation $f_n = 0$ (4) if

$$I_{n+1} - I_n = f_n \Lambda_n. \quad (15)$$

Although invariance of the integrals is implied by the staircase method, we will prove it directly by using properties of the multi-sums of products, Ψ and, at the same time, we obtain the integrating factors. From now on, for our convenience, we write u_i instead of u_{n+i} .

We denote $k = -(\lambda^2 + 1)/\lambda^2$, and we split the reduced \tilde{L} matrices as we did in [20]. We have

$$\tilde{L}_i = k u_{i+1} \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} 1 & u_{i+1} \\ \frac{1}{u_{i+1} + 1} & \frac{u_{i+1}}{u_{i+1} + 1} \end{pmatrix} = r_i (kH + s_i A_i^i), \quad (16)$$

where

$$r_i := u_{i+1}, s_i := \frac{1}{u_{i+1}(u_{i+1} + 1)}, H := \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \text{ and } A_j^i := \begin{pmatrix} a_i & a_i b_j \\ 1 & b_j \end{pmatrix},$$

with $a_i := u_{i+1} + 1$ and $b_i := u_{i+1}$. Therefore, applying Lemma 8 in [20], we obtain the following

$$\prod_{i=a}^{\hat{b}} \tilde{L}_i = \left(\sum_{r=0}^{b-a+1} X_r^{a,b} k^r \right) \prod_{i=a}^b r_i,$$

with

$$X_r^{a,b} = \Psi_{r-1}^{a+1, b-2} H A_a^{b-1} + \Psi_{r-1}^{a+2, b-1} A_{a+1}^b H + \Psi_{r-2}^{a+2, b-2} H + \Psi_r^{a+1, b-1} A_a^b, \quad (17)$$

where $c_i := s_i(a_{i-1} + b_i)$ and

$$\Psi_r^{a,b} := s_{a-1} \left(\sum_{a \leq i_1, i_1+1 < i_2, i_2+1 < \dots < i_{r-1}, i_{r-1}+1 < i_r \leq b} \prod_{j=1}^r f_{i_j} \right) \prod_{i=a}^{b+1} c_i, \quad (18)$$

with $f_i := s_{i+1}/(c_i c_{i+1})$.

By writing

$$(1+k)\tilde{M}_0^{-1} = kH + \begin{pmatrix} 0 & 1 \\ \frac{1}{(u_0+1)u_p} & \frac{-1}{(u_0+1)u_p} \end{pmatrix},$$

we obtain

$$(k+1)\text{Tr}(\tilde{\mathcal{L}}_0) = \sum_{r=0}^{\lfloor (p+1)/2 \rfloor} k^r I_r^p \quad (19)$$

where I_r^p is given as follows

$$I_r^p = \left(\frac{u_1 \Psi_{r-1}^{1,p-3}}{(u_0+1)u_p} + \frac{\Psi_{r-1}^{2,p-2}}{u_0+1} + \frac{\Psi_{r-2}^{2,p-3}}{u_p(u_0+1)} + \frac{(u_0+u_1+1)\Psi_r^{1,p-2}}{u_0+1} + \Psi_{r-1}^{1,p-2} \right) \prod_{i=1}^p u_i, \quad (20)$$

with

$$s_i = \frac{1}{u_{i+1}(u_{i+1}+1)}, \quad c_i = \frac{u_i+u_{i+1}+1}{u_{i+1}(u_{i+1}+1)}, \quad f_i = \frac{u_{i+1}(u_{i+1}+1)}{(u_i+u_{i+1}+1)(u_{i+1}+u_{i+2}+1)}.$$

We also get

$$\frac{(k+1)}{(-k)^{p+1}} \text{Det} \tilde{\mathcal{L}}_0 = \frac{u_1 u_2 \dots u_{p-1}}{(u_0+1)(u_1+1) \dots (u_p+1)} =: I_G, \quad (21)$$

which gives us the inverse of the integral (4.1) given in [9]. Thus, we have the following theorem.

Theorem 1. *Let $p > 1$ and $0 \leq 2r \leq p+1$. We have*

(i) I_r^p given by (20) is an integral of equation (2) with integrating factor

$$\Lambda_r^p = \left(\frac{\Psi_{r-1}^{2,p-2}}{(u_0+1)(u_{p+1}+1)} + \frac{\Psi_{r-2}^{2,p-3}}{u_p(u_0+1)(u_{p+1}+1)} + \frac{u_1 u_{p+1} \Psi_r^{1,p-1}}{(u_0+1)u_p} - \frac{u_1 \Psi_r^{1,p-2}}{(u_0+1)u_p} \right) \prod_{i=2}^p u_i \quad (22)$$

(ii) I_G given by (21) is also an integral of equation (2) with integrating factor

$$\Lambda_{I_G} = \frac{u_2 u_3 \dots u_{p-1}}{(u_0+1)(u_1+1) \dots (u_p+1)(u_{p+1}+1)}. \quad (23)$$

We prove this theorem directly using the following properties of Ψ given in [20]

$$\Psi_r^{n,m} = s_{m+1}(a_m + b_{m+1})\Psi_r^{n,m-1} + s_{m+1}\Psi_{r-1}^{n,m-2}, \quad (24)$$

$$\Psi_r^{n,m} = s_{n-1}(a_{n-1} + b_n)\Psi_r^{n+1,m} + s_{n-1}\Psi_{r-1}^{n+2,m}. \quad (25)$$

Proof. (i) We write

$$S(I_r^p) - I_r^p = \prod_{i=2}^p u_i (A + B), \quad (26)$$

where S is a shift operator, i.e $S(u_i) = u_{i+1}$ and

$$A = \left(\frac{u_2 \Psi_{r-1}^{2,p-2}}{u_1+1} + \frac{\Psi_{r-2}^{3,p-2}}{u_1+1} + u_{p+1} \Psi_{r-1}^{2,p-1} \right) - u_1 \left(\frac{\Psi_{r-1}^{2,p-2}}{u_0+1} + \frac{\Psi_{r-2}^{2,p-3}}{u_p(u_0+1)} + \Psi_{r-1}^{1,p-2} \right),$$

$$B = u_{p+1} \left(\frac{\Psi_{r-1}^{3,p-1}}{u_1+1} + \frac{(u_1+u_2+1)\Psi_r^{2,p-1}}{u_1+1} \right) - u_1 \left(\frac{u_1 \Psi_{r-1}^{1,p-3}}{(u_0+1)u_p} + \frac{(u_0+u_1+1)\Psi_r^{1,p-2}}{u_0+1} \right).$$

Using properties (24) and (25), we have

$$A = \left(\frac{u_2 \Psi_{r-1}^{2,p-2}}{(u_1+1)} + \frac{\Psi_{r-2}^{3,p-2}}{(u_1+1)} + \frac{(u_p+u_{p+1}+1)\Psi_{r-1}^{2,p-2}}{u_{p+1}+1} + \frac{\Psi_{r-2}^{2,p-3}}{u_{p+1}+1} \right)$$

$$- u_1 \left(\frac{\Psi_{r-1}^{2,p-2}}{u_0+1} + \frac{\Psi_{r-2}^{2,p-3}}{u_p(u_0+1)} + \frac{(u_1+u_2+1)\Psi_{r-1}^{2,p-2}}{u_1(u_1+1)} + \frac{\Psi_{r-2}^{3,p-2}}{u_1(u_1+1)} \right)$$

$$= \left(u_p(u_0+1) - u_1(u_{p+1}+1) \right) \left(\frac{\Psi_{r-1}^{2,p-2}}{(u_0+1)(u_{p+1}+1)} + \frac{\Psi_{r-2}^{2,p-3}}{u_p(u_0+1)(u_{p+1}+1)} \right).$$

Similarly, we get

$$B = \left(u_p(u_0 + 1) - u_1(u_{p+1} + 1) \right) \left(\frac{u_1 u_{p+1} \Psi_r^{1,p-1}}{(u_0 + 1)u_p} - \frac{u_1 \Psi_r^{1,p-2}}{(u_0 + 1)u_p} \right).$$

Therefore, we obtain an integrating factor as given in (22).

(ii) By direct calculation, one obtains the integrating factor Λ_{IG} given by (23). □

5. Integrals of the p^{th} order Lyness equation

In this section, we construct integrals of the original Lyness equation from the corresponding derivative Lyness equation. We also show that the integrals given by Gao et al in [8] can be derived as a special cases from integrals obtained from our closed-form expressions.

Let us assume that $I(u_0, u_1, \dots, u_p)$ is an integral of the derivative Lyness equation (2). This means

$$I(u_0, u_1, \dots, u_p) = I(u_1, u_2, \dots, u_p, \frac{u_0 u_p + u_p - u_1}{u_1}). \quad (27)$$

Now we define

$$Q(u_0, u_1, \dots, u_{p-1}) = I(u_0, u_1, \dots, u_{p-1}, \frac{a + u_1 + u_2 + \dots + u_{p-1}}{u_0}). \quad (28)$$

Lemma 2. $Q(u_0, u_1, \dots, u_{p-1})$ is an integral of the Lyness equation (1), i.e

$$Q(u_0, u_1, \dots, u_{p-1}) = Q(u_1, u_2, \dots, u_{p-1}, \frac{a + u_1 + u_2 + \dots + u_{p-1}}{u_0}). \quad (29)$$

Proof. It can be seen that

$$\begin{aligned} LHS(29) &= I(u_0, u_1, \dots, u_{p-1}, \frac{a + u_1 + u_2 + \dots + u_{p-1}}{u_0}) \\ &= I(u_1, u_2, \dots, u_{p-1}, \frac{a + u_1 + u_2 + \dots + u_{p-1}}{u_0}, \frac{u_0 \frac{a + u_1 + u_2 + \dots + u_{p-1}}{u_0} + \frac{a + u_1 + u_2 + \dots + u_{p-1}}{u_0} - u_1}{u_1}) \\ &= I(u_1, u_2, \dots, u_{p-1}, \frac{a + u_1 + u_2 + \dots + u_{p-1}}{u_0}, \frac{a + u_2 + \dots + u_{p-1} + \frac{a + u_1 + u_2 + \dots + u_{p-1}}{u_0}}{u_1}) \\ &= Q(u_1, u_2, \dots, u_{p-1}, \frac{a + u_1 + u_2 + \dots + u_{p-1}}{u_0}) \end{aligned}$$

This means that $Q(u_0, u_1, \dots, u_{p-1})$ is an integral for the Lyness equation. □

Using this theorem, it is easy to see that the first invariant in [8] is derived from I_G . For the others, by writing Ψ explicitly in terms of the variables u , one can show that the second and third invariants in [8] are equivalent to I_0^p/I_G and $I_{[(p+1)/2]}^p/I_G$ respectively.

By using Theorem 1 and Lemma 2, we obtain closed-form expressions for integrals of the p^{th} order Lyness equation. In the following corollary of Theorem 1 we have exploited property (24) to make the dependence of I_r^p of its last argument explicit.

Corollary 3. We denote $T := a + u_0 + u_1 + \dots + u_{p-1}$. Let $p > 2$ and $0 \leq 2r \leq p + 1$. Then integrals of the p^{th} order Lyness equation are given by

$$Q_{I_G} = \frac{u_0 u_1 \dots u_{p-1}}{(u_0 + 1)(u_1 + 1) \dots (u_{p-1} + 1)T}, \quad (30)$$

$$Q_r^p = \left(\frac{u_1 \Psi_{r-1}^{1,p-3}}{u_0 + 1} + \frac{(T + u_0 u_{p-1}) \Psi_{r-1}^{2,p-3}}{T(u_0 + 1)} + \frac{u_0 \Psi_{r-2}^{2,p-4}}{T(u_0 + 1)} + \frac{\Psi_{r-2}^{2,p-3}}{u_0 + 1} + \frac{u_0(u_0 + u_1 + 1) \Psi_{r-1}^{1,p-4}}{T(u_0 + 1)} \right. \\ \left. + \frac{(u_0 + u_1 + 1)(T + u_0 u_{p-1}) \Psi_r^{1,p-3}}{T(u_0 + 1)} + \frac{(T + u_0 u_{p-1}) \Psi_{r-2}^{1,p-3}}{T} + \frac{u_0 \Psi_{r-2}^{1,p-4}}{T} \right) \prod_{i=1}^{p-1} u_i. \quad (31)$$

6. Some relations between the integrals

In this section, we first give some relations between the integrals I_r^p of the derivative Lyness equation given by (20), and the integral I_G given by (21). This will lead to some relations between the integrals of the Lyness equation.

6.1. Some relations between the integrals of the derivative Lyness equation

We denote the monodromy matrix obtained by using the Lax pair $(L_{l,m}/\lambda, M_{l,m})/\lambda$ by \mathcal{L}_0 where (L, M) are given by (11). We have

$$\mathcal{L}_0 = \left(\frac{M_0}{\lambda} \right)^{-1} \frac{L_{p-1}}{\lambda} \frac{L_{p-2}}{\lambda} \dots \frac{L_0}{\lambda}.$$

It is easy to see that $\tilde{\mathcal{L}}_0 = G_0^{-1} \mathcal{L}_0 G_0$, where G_0 is given by (13). This implies

$$\text{Tr}(\tilde{\mathcal{L}}_0) = \text{Tr} \mathcal{L}_0. \quad (32)$$

Recall that integrals I_r^p are obtained by expanding $(k + 1)\text{Tr} \tilde{\mathcal{L}}_0$ in powers of $k = -1 - 1/\lambda^2$, see equation (19). Now we expand $(k + 1)\text{Tr} \mathcal{L}_0$ in powers of $1/\lambda$. Equating the lowest coefficient and the coefficient for $1/\lambda^2$ we obtain the following relations.

Proposition 4. Let $p \geq 2$, let I_r^p and I_G be given in (20) and (21). Then we have

$$\sum_{r=0}^{\lfloor (p+1)/2 \rfloor} (-1)^r I_r^p = -I_G, \quad (33)$$

$$\sum_{r=1}^{\lfloor (p+1)/2 \rfloor} (-1)^r r I_r^p + 1 = (u_0 u_p - u_1 - u_2 - \dots - u_{p-1} - p) I_G. \quad (34)$$

A proof of this proposition will be given in the Appendix.

Remark 5. Note that $u_0 u_p - u_1 - u_2 - \dots - u_{p-1}$ is also an integral of the derivative Lyness equation with integrating factor 1.

6.2. Some relations between the integrals of the Lyness equation

Using Proposition 4, we obtain the following linear relations between the integrals of the Lyness equation.

Corollary 6. *Let $p \geq 3$, let Q_r^p, Q_{IG} be given by (31) and (30). Then we have*

$$\sum_{r=0}^{\lfloor (p+1)/2 \rfloor} (-1)^r Q_r^p = -Q_{IG}, \quad (35)$$

$$\sum_{r=0}^{\lfloor (p+1)/2 \rfloor} (-1)^r (p - a - r) Q_r^p = 1. \quad (36)$$

7. Conclusion

We have presented closed-form expressions for integrals of the p^{th} order Lyness equation and its derivative equation. Using Maple, for $p \leq 11$, for random values u_i we have found that the set of integrals $\{I_0^p, I_1^p, \dots, I_{\lfloor (p+1)/2 \rfloor}^p\}$ of the derivative Lyness equation is functionally independent. Taking particular values for the variables we are able to check orders that are a bit higher. We found functional independence for $p \leq 15$ when $u_0 = u_1 = \dots = u_{p-1} = 1$ and $u_p = 2$. For the p^{th} order Lyness equation the set $\{Q_0^p, Q_1^p, \dots, Q_{\lfloor (p+1)/2 \rfloor}^p\}$ is not functionally independent due to the linear relation (36). However, the rank of the Jacobian matrix seems to be $R = \lfloor (p+1)/2 \rfloor$ for $p \leq 11$ and for random value a and u_i . In particular, this is the case for $p \leq 15$ and for random value of a when $u_0 = u_1 = \dots = u_{p-1} = 1$. In fact, for all $p \leq 15$ the determinant of $2 \frac{a+p}{a+p-2}$ times the upperleft $R \times R$ part of the Jacobian is

$$\pm \frac{3^{q_p-1}}{2^{g_p+1}} \frac{a - \lfloor p/2 \rfloor}{a+p}, \quad (37)$$

where the powers of 3 are given by quarter squares $q_p = \lfloor p^2/4 \rfloor$ and the powers of 2 are generalised pentagonal numbers, cf [19],

$$g_p = \lfloor \frac{p}{2} \rfloor \left(3 \lfloor \frac{p}{2} \rfloor - (-1)^p \right) / 2.$$

We conjecture the formula (37) to be true for any p . The existence of $\lfloor (p+1)/2 \rfloor$ functionally independent integrals in the set our integrals for the Lyness equation would be enough for completely integrability in the sense of Liouville-Arnold.

Another problem is to see if the Lyness equation has a symplectic structure or not. One way to approach this problem is using a Lagrangian to derive the symplectic structures [4]. We have found a Lagrangian for a double copy of equation (2). However, to construct symplectic structures for the Lyness mapping itself is still an open problem.

Acknowledgments

This research has been funded by the Australian Research Council through the Centre of Excellence for Mathematics and Statistics of Complex Systems. Dinh T. Tran acknowledges the support of two scholarships, one from La Trobe University and the other from the Endeavour IPRS programme.

Appendix A. Proof of proposition 4

We write

$$L_i/\lambda = \begin{pmatrix} 1 & 0 \\ 0 & \frac{u_{i+1}}{u_{i+1}+1} \end{pmatrix} + \frac{1}{\lambda} \begin{pmatrix} 0 & -u_{i+1} \\ \frac{1}{u_{i+1}+1} & 0 \end{pmatrix}, \quad (\text{A.1})$$

and

$$(k+1)(M_0/\lambda)^{-1} = \begin{pmatrix} 0 & 0 \\ 0 & \frac{-1}{(u_0+1)u_p} \end{pmatrix} + \frac{1}{\lambda} \begin{pmatrix} 0 & -1 \\ \frac{1}{(u_0+1)u_p} & 0 \end{pmatrix}. \quad (\text{A.2})$$

Equalling the lowest coefficient of the following

$$(k+1)\text{Tr}(\tilde{\mathcal{L}}_0) = (k+1)\text{Tr}\mathcal{L}_0,$$

we obtain the first relation (33). Now equalling the coefficient of $1/\lambda^2$, we obtain

$$\begin{aligned} \sum_{r=1}^{\lfloor (p+1)/2 \rfloor} (-1)^r r I_r^p &= \left(\sum_{1 \leq r < s \leq p} \prod_{i=1}^r u_i \prod_{i=r}^{s-1} (1+u_i) \prod_{i=s+1}^p u_i - \sum_{r=1}^p \left(u_p \prod_{i=r+1}^p u_i \prod_{i=0}^{r-1} (1+u_i) \right. \right. \\ &\quad \left. \left. + \prod_{i=1}^r u_i \prod_{i=r}^p (1+u_i) \right) \right) \frac{1}{u_p(u_0+1)(u_1+1)\dots(u_p+1)}. \end{aligned} \quad (\text{A.3})$$

In this proof, we use the following identity with $m < n$

$$\prod_{i=m}^n (1+u_i) - \sum_{m < s \leq n} \prod_{i=m}^{s-1} (1+u_i) \prod_{i=s+1}^n u_i = (1+u_m) \prod_{i=m+1}^n u_i. \quad (\text{A.4})$$

This identity is proved by expanding its left hand side. We have

$$\begin{aligned} LHS(\text{A.4}) &= \left(\prod_{i=m}^n (1+u_i) - \prod_{i=m}^{n-1} (1+u_i) \right) - \sum_{m < s \leq n-1} \prod_{i=m}^{s-1} (1+u_i) \prod_{i=s+1}^n u_i \\ &= \left(\prod_{i=m}^{n-1} (1+u_i) - \prod_{i=m}^{n-2} (1+u_i) \right) u_n - \sum_{m < s \leq n-2} \prod_{i=m}^{s-1} (1+u_i) \prod_{i=s+1}^n u_i \\ &\quad \vdots \\ &= (1+u_m) \prod_{i=m+1}^n u_i = RHS(\text{A.4}). \end{aligned}$$

We write

$$\sum_{r=1}^{\lfloor (p+1)/2 \rfloor} (-1)^r r I_r + 1 = \frac{P_1}{P_2},$$

where

$$\begin{aligned} P_1 &= u_p \prod_{i=0}^p (u_i+1) + \sum_{1 \leq r < s \leq p} \prod_{i=1}^r u_i \prod_{i=r}^{s-1} (1+u_i) \prod_{i=s+1}^p u_i - \sum_{r=1}^p u_p \prod_{i=r+1}^p u_i \prod_{i=0}^{r-1} (1+u_i) \\ &\quad + \prod_{i=1}^r u_i \prod_{i=r}^p (1+u_i), \end{aligned}$$

$$P_2 = u_p \prod_{i=0}^p (u_i+1).$$

Thus, we only need to show that

$$P_1 = (u_0 u_p - u_1 - u_2 - \dots - u_{p-1} - p) \prod_{i=1}^p u_i.$$

We have

$$\begin{aligned} P_1 &= \left(u_p \prod_{i=0}^p (u_i + 1) - \sum_{r=1}^p u_p \prod_{i=r+1}^p u_i \prod_{i=0}^{r-1} (1 + u_i) \right) + \sum_{1 \leq r < s \leq p-1} \prod_{i=1}^r u_i \prod_{i=r}^{s-1} (1 + u_i) \prod_{i=s+1}^p u_i \\ &\quad + \sum_{1 \leq r \leq p-1} \prod_{i=1}^r u_i \prod_{i=r}^{p-1} (1 + u_i) - \sum_{r=1}^p \prod_{i=1}^r u_i \prod_{i=r}^p (1 + u_i) \\ &= u_p (1 + u_0) \prod_{i=1}^p u_i + \sum_{1 \leq r < s \leq p-1} \prod_{i=1}^r u_i \prod_{i=r}^{s-1} (1 + u_i) \prod_{i=s+1}^p u_i - \sum_{1 \leq r \leq p-1} \prod_{i=1}^r u_i \prod_{i=r}^{p-1} (1 + u_i) u_p \\ &\quad - (1 + u_p) \prod_{i=1}^p u_i \\ &= (u_0 u_p - 1) \prod_{i=1}^p u_i + \sum_{r=1}^{p-1} \prod_{i=1}^r u_i \left(\sum_{r < s \leq p-1} \prod_{i=r}^{s-1} (1 + u_i) \prod_{i=s+1}^{p-1} u_i - \prod_{i=r}^{p-1} (1 + u_i) \right) u_p \\ &= (u_0 u_p - 1) \prod_{i=1}^p u_i - \sum_{r=1}^{p-1} (1 + u_r) \prod_{i=1}^p u_i = (u_0 u_p - p - u_1 - u_2 - \dots - u_{p-1}) \prod_{i=1}^p u_i. \end{aligned}$$

This proves our statement.

References

- [1] Adler V, Bobenko A and Suris Y 2003 Classification of Integrable Equations on quad-graphs. The Consistency Approach *Commun. Math. Phys.* **233** 513-43
- [2] Bastien G and Rogalski M 2004 Global behavior of the solutions of Lyness' difference equation $u_{n+2}u_n = u_{n+1} + a$ *J. Difference Equ. Appl.* **10** 977-1003
- [3] Bastien G and Rogalski M 2009 Results and conjectures about order q Lyness' difference equation $u_{n+q}u_n = a + u_{n+q-1} + \dots + u_{n+1}$ in \mathbb{R}_*^+ , with a particular study of the case $q = 3$ *Adv. Difference Equ.* **2009** Article ID 134749 (36pp)
- [4] Bruschi M, Ragnisco O, Santini P M and Gui-Zhang T 1991 Integrable symplectic maps *Physica D* **49** 273-94
- [5] Cima A, Gasull A and Mañosa 2007 Dynamics of the third order Lyness difference equation *J. Difference Equ. Appl.* **13** 855-84.
- [6] Cima A, Gasull A and Mañosa V 2008 Some properties of the k -dimensional Lyness' map *J. Phys. A: Math. Theor.* **41** 285205-22
- [7] Darwen C and Patula W T 1998 Properties of a certain Lyness equation *J. Math. Anal. Appl.* **218** 458-78
- [8] Gao M, Kato Y and Ito M 2004 Some invariants for k th-order Lyness equation *Appl. Math. Lett.* **17** 1183-89
- [9] Grammaticos B, Ramani A and Tamizhmani T 2009 Investigating the integrability of the Lyness mapping *J. Phys. A: Math. Theor.* **42** 454009-15
- [10] Kocic V L, Ladas G and Rodriguez I W 1993 On rational recursive sequences *J. Math. Ana. App.* **173** 127-57

- [11] Kulenović M R S 2000 Invariants and related Liapunov functions for difference equations *Appl. Math. Lett* **13** 1–8
- [12] Ladas, G 1995 Open problems and conjectures *J. Difference Equ. Appl.* **1** 209–14
- [13] Levi D and Yamilov R I 2009 The generalized symmetry method for discrete equations *J. Phys. A: Math. Theor.* **42** 454012–29
- [14] Lyness R C 1942 Note 1581 *The Mathematical Gazette* **26** 62
- [15] Papageorgiou V G, Nijhoff F W and Capel H W 1990 Integrable mappings and nonlinear integrable lattice equations *Phys. Lett. A* **147** 106–14
- [16] Quispel G R W, Capel H W, Papageorgiou V G and Nijhoff F W 1991 Integrable mappings derived from soliton equations *Physica A* **173** 243–66
- [17] Quispel G R W, Capel H W and Roberts J A G 2005 Duality for discrete integrable systems, *J. Phys. A: Math. Gen.* **38** 3965–80
- [18] Rojas O, van der Kamp P H and Quispel G R W Lax representations for integrable maps, *in preparation*
- [19] Sloane N J A, The On-Line Encyclopedia of Integer Sequences, published electronically at www.research.att.com/njas/sequences/
- [20] Tran D T, van der Kamp P H and Quispel G R W 2009 Closed-form expressions for integrals of traveling wave reductions of integrable lattice equations *J. Phys. A: Math. Theor.* **42** 225201–20
- [21] van der Kamp P H 2009 Initial value problems for lattice equations *J. Phys. A: Math. Theor.* **42** 404019–34
- [22] van der Kamp P H and Quispel G R W The staircase method *in preparation*
- [23] Veselov A P 1991 Integrable maps, *Russian Math. Surveys* **46** 1–51