



References for depression intervention tables

Appleby, L., Warner, R., Whitton, A., & Faragher, B. A controlled study of fluoxetine and cognitive-behavioural counselling in the treatment of postnatal depression. *British Medical Journal*, 1997, 314, 932-36.

Armstrong, K.L., Fraser, J.A., Dadds, M.R., & Morris, J. A randomized controlled trial of nurse home visiting to vulnerable families with newborns. *Journal of Paediatrics and Child Health* , 1999, 35, 237-44.

Armstrong, K.L., Fraser, J.A., Dadds, M.R., & Morris, J. Promoting secure attachment, maternal mood and child health in a vulnerable population: a randomised controlled trial. *Journal of Paediatrics and Child Health* , 2000, 36, 555-62.

Biro, M.A., Waldenström, U., & Pannifex, J. Team midwifery care in a tertiary level obstetric service: a randomized controlled trial. *Birth*, 2000, 27, 174-6.

Biro, M.A., Pannifex, J., Tippett, C., Merriel, R., Brown, S., & Waldenström, U. Colative Pregnancy Care/Team Midwifery Project. Final Evaluation Report, Commonwealth Birthing Services Program – Victoria, 1999, pp 38.

Brugha, T.S., Wheatley, S., Taub, N.A. et al. Pragmatic randomised trial of an antenatal intervention to prevent post-natal depression by reducing psychosocial risk factors. *Psychological Medicine* 2000, 30,1273-81.

Brugha, T.S., Sharp, H.M., Cooper, S.A. et al. The Leicester 500 Project. Social support and the development of postnatal depressive symptoms, a prospective cohort survey. *Psychological Medicine*, 1998, 28, 63-79.

Buist, A., Westley, D., & Hill, C. Antenatal prevention of postnatal depression. *Archives of Women's Mental Health* 1998, 1, 1-7.

Chabrol, H., Teissedre, F., Saint-Jean, M. et al. Dépistage, prévention et traitement des dépressions du post-partum: une étude contrôlée chez 859 sujets. *L'Encéphale*, 2002, XXVIII, 65-70.

Chabrol, H., Teissedre, F., Saint-Jean, M. et al. Prevention and treatment of postpartum depression. A controlled randomized study on women at risk. *Psychological Medicine*, 2002, 32, 1039-47.

Chen, C-H., Tseng, Y-F., Chou, F-H., & Wang, S-Y. Effects of group support intervention in postnatally distressed women. A controlled study in Taiwan. *Journal of Psychosomatic Research*, 2000, 49, 395-99.

Cooper, P., & Murray, L. The impact of psychological treatments of postpartum depression on maternal mood and infant development. In: Murray, L., & Cooper, P. (eds.), *Postpartum depression and child development*. New York: the Guildford Press, 1997, 201-220.

Cooper, P., Murray, L., Wilson, A., Romaniuk, H. Controlled trial of the short- and long-term effect of psychological treatment of post-partum depression. I impact on maternal mood. *British Journal of Psychiatry* 2003, 182, 412-419.

Dennis, C.-L. The effect of peer support on postpartum depression. A pilot randomized controlled trial. *Canadian Journal of Psychiatry* 2003, 48, 115-124.

Elliott, S.A., Leverton, T.J., Sanjack, M., et al. Promoting mental health after childbirth: a controlled trial of primary prevention of postnatal depression. *British Journal of Clinical Psychology*, 2000, 39, 223-41.

Field, T., Grizzle, N., Scafidi, F., Schanberg, S. Massage and relaxation therapies' effects on depressed adolescent mothers. *Adolescence* 1996, 31, 903-11.

Fleming, A.S., Klein, E., & Corter, C. The effects of a social support group on depression, maternal attitudes and behaviour in new mothers. *Journal of Child Psychology and Psychiatry* 1992, 33, 685-98.

Gordon, N.P., Walton, D., McAdam, E. et al. Effects of providing hospital-based doulas in Health Maintenance Organization hospitals. *Obstetrics and Gynecology* , 1999, 93, 422-426.

Gordon, R.E., Gordon, K.K., & Englewood, N. Social factors in the prediction and treatment of emotional disorders of pregnancy. *American Journal of Obstetrics and Gynecology*, 1959, 77, 1074-83.

Gordon, R.E., & Gordon, K.K. Social factors in prevention of postpartum emotional problems. *Obstetrics and Gynecology*, 1960, 15, 433-37.

Gordon, R.E., Kapostins, E.E., & Gordon, K.K. Factors in postpartum emotional adjustments. *Obstetrics and Gynecology*, 1965, 25, 158-66.

Gunn, J., Lumley, J., Chondros, P., & Young, D. Does an early postnatal check-up improve maternal health: results from a randomised trial in Australian general practice. *British Journal of Obstetrics and Gynaecology*, 1998, 105, 991-97.

Hayes, B.A., Muller, R., & Bradley, B.S. Perinatal depression: a randomised controlled trial of an antenatal education intervention for primiparas. *Birth*, 2001, 28, 28-35.

Heinicke, C.M., Fineman, N.R., Ruth, G. et al. Relationship-based intervention with at-risk mothers: outcome in the first year of life. *Infant Mental Health Journal*, 1999, 20, 349-74.

Hiscock, H., & Wake, M. Randomised controlled trial of behavioural infant sleep intervention to improve infant sleep and maternal mood. *British Medical Journal* 2002, 324, 1062-65.

Hodnett, E.D., Lowe, N.K., Hannah, M.E., et al. Effectiveness of nurses as providers of birth support in North American Hospitals. A randomized controlled trial. *JAMA*, 2002, 288, 1373-81.

Hoffman, Y. The effects of a supportive intervention during and delivery on the postpartum psychological adaptation of first-time mothers. PhD thesis. Case Western Reserve University, 1992.

Holden, J.M., Sagovsky, R., & Cox, J.L. Counselling in a general practice setting: controlled study of health visitor intervention in treatment of postnatal depression. *British Medical Journal*, 1989, 298, 223-26.

Honey, K. L, Bennett, P., Morgan, M. A brief psycho-educational group intervention for postnatal depression. *British Journal of Clinical Psychology* 2002, 41, 405-409.

Horowitz, J.A., Bell, M., Trybulski J., et al. Promoting responsiveness between mothers with depressive symptoms and their infants. *Journal of Nursing Scholarship*, 2001, 33, 323-39.

Lavender, T., & Walkinshaw, S.A. Can midwives reduce postpartum psychological morbidity? A randomised trial. *Birth*, 1998, 25, 215-19.

Leverton, T.J., & Elliott, S.A. Transition to parenthood groups: a preventive intervention for postnatal depression? In Van Hall, E.V., & Everaerd, W. (eds.), *The Free Woman: Women's Health in the 1990s*. Carnforth: The Parthenon Publishing Group, 1989, 479-86. [see Elliott].

Marks, M. N., Siddle, K., Warwick, C. Can we prevent postnatal depression? A randomised controlled trial to assess the effect of continuity of midwife care on rates of postnatal depression in high risk women. *Journal of Maternal-Fetal Medicine* 2003, 13, 119-127.

MacArthur, C., Winter, H., Bick, D. et al. Effects of redesigned community postnatal care on women's health 4 months after birth: a cluster randomised trial. *Lancet* 2002, 359, 378-85.

MacArthur, C., Winter, H., Bick, D. et al. Re-designing postnatal care: a randomised control trial of protocol-based midwifery-led care focussed on women's physical and psychological health needs. *Health Technology Assessment* 2003, 7, 37.

Misri, S., Kostaras, X., Fox, D., & Kostaras, D. The impact of partner support in the treatment of postpartum depression. *Canadian Journal of Psychiatry*, 2000, 45, 554-58.

Morrell, C.J., Spiby, H., Stewart, P., Walters, S., & Morgan, A. Costs and effectiveness of community postnatal support workers: randomised controlled trial. *British Medical Journal*, 2000, 321, 593-98.

Morrell, C.J., Spiby, H., Stewart, P., Walters, S., & Morgan, A. Costs and benefits of community postnatal support workers: a randomised controlled trial. *Health Technology Assessment* , 2000, 4, 6.

Murray, L., Cooper, P., Wilson, A., Romaniuk, H. Controlled trial of the short- and long-term effect of psychological treatment of postpartum depression. 2. Impact on the mother-child relationship. *British Journal of Psychiatry* 2003, 182, 420-427. [see Cooper].

Nikodem, V.C., Nolte, A.G., Wolman, W., Gulmezoglu, A.M., & Hofmeyr, G.J. Companionship by a lay labor supporter to modify the clinical birth environment: long-term effects on mother and child. *Curationis*, 1998, 21, 8-12.

O'Hara, M.W., Stuart, S., Gorman, L.L., & Wenzel, A. Efficacy of interpersonal therapy for postpartum depression. *Archives of General Psychiatry* 2000, 57, 1039-45.

Onazawa, K., Glover, V., Adams, D., Modi, N., & Kumar, R.C. Infant massage improves mother-infant interaction for mothers with postnatal depression. *Journal of Affective Disorders*, 2001, 63, 201-07.

Priest, S., Henderson, J., R., Evans, S.F., Hagan, R.. Stress debriefing after childbirth: a randomised controlled trial. *Medical Journal of Australia* 2003, 178, 542-545.

Reid, M., Glazener, C., Murray, G., & Taylor, G.S. A two-centred pragmatic randomised controlled trial of two interventions of postnatal support. *British Journal of Obstetrics and Gynaecology*, 2002, 109, 1164-1170.

Shields, N., Reid, M., Cheyne, H. et al. Impact of midwife-managed care in the postnatal period: an exploration of psychosocial outcomes. *Journal of Reproductive and Infant Psychology*, 1997, 15 , 91-108.

Small, R., Lumley, J., Donohue, L., Potter, A., & Waldenström. Randomised controlled trial of midwife led debriefing to reduce maternal depression after operative childbirth. *British Medical Journal*, 2000, 321, 1043-47.

Spinelli, M., Endicott, J. Controlled clinical trial of interpersonal psychotherapy versus parenting education program for depressed pregnant women. *American Journal of Psychiatry* 2003,160, 555-562.

Stamp, G.E., Williams, A.S., & Crowther, C. Evaluation of antenatal and postnatal support to overcome postnatal depression. A randomised controlled trial. *Birth*, 1995, 22, 138-43.

Turnbull, D., Holmes, A., Shields, N., et al. Randomised, controlled trial of efficacy of midwife-managed care. *Lancet* 1996; 348: 213-18.

Waldenström, U., Brennecke, S., Brown, S et al.. Team Midwife Care. Final Report to the Commonwealth Birthing Services Program – Victoria, 2000, pp 30.

Waldenström, U., McLachlan, H., Forster, D., Brennecke, S., & Brown, S. Team midwife care: maternal and infant outcomes. *Australian and New Zealand Journal of Obstetrics and Gynaecology*, 2001, 41, 257-64.

Webster, J., Linnane, J., Roberts, J., Starrenburg, S., Hinson, J., Dibley, L. Identify, Educate and Alert (IDEA) trial: an intervention to reduce postnatal depression. *BJOG* 2003, 110, 842-846.

Wheatley, S.L., & Brugha, T.S. 'Just because I like it doesn't mean it has to work': personal experiences of an antenatal psychosocial intervention designed to prevent postnatal depression. *International Journal of Mental Health Promotion*, 1999, 1, 26-31. [see Brugha].

Wheatley, S.L., Culverwell, A., Brugha, T.S., & Shapiro, D.A. Preparing for Parenthood: background and development of a risk modifying intervention to prevent postnatal depression. *Archives of Women's Mental Health*, 2000, 3, 81-90. [see Brugha]

Wickberg, B., & Hwang, C.P. Counselling of postnatal depression: A controlled study on a population based Swedish sample. *Journal of Affective Disorders*, 1996, 39, 209-16.

Wolman, W.L., Chalmers, B., Hofmeyr, J., & Nikodem, V.C. Postpartum depression and companionship in the clinical birth environment. A randomised controlled study. *American Journal of Obstetrics and Gynecology* 1993, 168, 1388-93.

Zlotnick, C., Johnson, S.L., Miller, I., Pearlstein, T., & Howard, M. Postpartum depression in women receiving public assistance: pilot study of an interpersonal-therapy-oriented group intervention. *American Journal of Psychiatry* 2001, 158, 638-40.